Brain Research, 561 (1991) 358–359
Elsevier

Author Index

Akaike, N., see Shirasaki, T., 77
Akiyama, Y., Ito, A., Koshimura, K., Ohue, T., Yamagata, S., Miwa, S. and Kikuchi, H., Effects of transient forebrain ischemia and reperfusion on function of dopaminergic neurons and dopamine reuptake in vivo in rat striatum, 120
Appenteng, K., see Saha, S., 128
Arnold, A.P., see Schlinger, B.A., 99

Ballinger, M.L., see Krause, T.L., 350
Batten, T.F.C., see Saha, S., 128
Becker, C.-M., see Naas, E., 139
Becker, D.P., see Yoshino, A., 106
Beitner-Johnson, D., Guitart, X. and Nestler, E.J., Dopaminergic brain-reward regions of Lewis and Fischer rats display different levels of tyrosine hydroxylase and other morphine- and cocaine-regulated phosphoproteins, 147
Betz, H., see Naas, E., 139
Bicknell, R.J., see Oldfield, B.J., 151
Bilkey, D.K., see Yamada, N., 324
Bitran, D., Hilvers, R.J. and Kellogg, C.K., Anxiolytic effects of 3α-hydroxy-5α[β]-pregnan-20-one: endogenous metabolites of progesterone that are active at the $GABA_A$ receptor, 157
Bittner, G.D., see Krause, T.L., 350
Bohn, M.C., see Cunningham, L.A., 192
Breedlove, S.M., see Leslie, M., 269
Breese, G.R., see Duncan, G.E., 84
Bridges, R.J., Hatalski, C., Shim, S.N. and Nunn, P.B., Gliotoxic properties of the *Lathyrus* excitotoxin β-*N*-oxalyl-L-α,β-diaminopropionic acid (β-L-ODAP), 262
Brody, M.J., see Meller, S.T., 61

Campbell, J.N., see Meyer, R.A., 252
Cerne, R., see Gerber, R., 236
Chalmers, D.T. and Watson, S.J., Comparative anatomical distribution of 5-HT_{1A} receptor mRNA and 5-HT_{1A} binding in rat brain — a combined in situ hybridisation/in vitro receptor autoradiographic study, 51
Chapin, J.K., see Nicolelis, M.A.L., 344
Chiu, A.T., see DeNoble, V.J., 230
Cleary, J., see Jewett, D.C., 165
Cler, J.A., see Rao, T.S., 43
Cohen, R.H., see Meyer, R.A., 252
Connor, J.A., see Hyndman, A.G., 318
Contreras, P.C., see Rao, T.S., 43
Cotman, C.W., see Koh, J.-y., 338
Crawford, C., see Iwamoto, N., 177
Cunningham, L.A., Hansen, J.T., Short, M.P. and Bohn, M.C., The use of genetically altered astrocytes to provide nerve growth factor to adrenal chromaffin cells grafted into the striatum, 192

Davis, K.D., see Meyer, R.A., 252
Deng, M.H., Lynch, H.J. and Wurtman, R.J., Melatonin and its precursors in Y79 human retinoblastoma cells: effect of sodium butyrate, 274
DeNoble, K.F., see DeNoble, V.J., 230
DeNoble, V.J., DeNoble, K.F., Spencer, K.R., Chiu, A.T., Wong, P.C. and Timmermans, P.B.M.W.M., Nonpeptide angiotensin II receptor antagonist and angiotensin-converting enzyme inhibitor: effect on a renin-induced deficit of a passive avoidance response in rats, 230
Dickson, K.L., see Seckl, J.R., 332
Dilworth, V.M., see Rao, T.S., 43
Dufresne, C., see McIntyre, D.C., 279
Duncan, G.E., Little, K.Y., Koplas, P.A., Kirkman, J.A., Breese, G.R. and Stumpf, W.E., β-Adrenergic receptor distribution in human and rat hippocampal formation: marked species differences, 84

Emmett, M.R., see Rao, T.S., 43
Emori, K., see Minabe, Y., 162
Emson, P.C., see Iwamoto, N., 177

Farley, L.G., see Silver, W.L., 212
Finger, T.E., see Silver, W.L., 212
Fink, G., see Seckl, J.R., 332
Forger, N.G., see Leslie, M., 269
Frusciante, M.J., see Nussenzveig, I.Z., 285

Gebhart, G.F., see Meller, S.T., 61
Gerber, G., Cerne, R. and Randić, M., Participation of excitatory amino acid receptors in the slow excitatory synaptic transmission in rat spinal dorsal horn, 236
Gibson, M.J., see Saitoh, Y., 11
Gnahn, H., see Naas, E., 139
Gray, N.M., see Rao, T.S., 43
Guitart, X., see Beitner-Johnson, D., 147

Haji, A., see Takeda, R., 307
Handa, N., see Kitagawa, K., 203
Hansen, J.T., see Cunningham, L.A., 192
Hata, R., see Kitagawa, K., 203
Hatalski, C., see Bridges, R.J., 262
Hilvers, R.J., see Bitran, D., 157
Hockberger, P.E., see Hyndman, A.G., 318
Hoebel, B.G., see Rada, P., 354
Horie, H., Pamplin, P.J. and Yokota, T., Inhibition of nociceptive neurons in the shell region of nucleus ventralis posterolateralis following conditioning stimulation of the periaqueductal grey of the cat. Evidence for an ascending inhibitory pathway, 35
Hovda, D.A., see Yoshino, A., 106
Hutchins, B., Evidence for a direct retinal projection to the anterior pretectal nucleus in the cat, 169
Hyndman, A.G., Hockberger, P.E., Zeevalk, G.D. and Connor, J.A., Transferrin can alter physiological properties of retinal neurons, 318

Ito, A., see Akiyama, Y., 120
Iwamoto, N., Thangnipon, W., Crawford, C. and Emson, P.C., Localization of calpain immunoreactivity in senile plaques and in neurones undergoing neurofibrillary degeneration in Alzheimer's disease, 177
Iyengar, S., see Rao, T.S., 43

Javitt, D.C., see Nussenzveig, I.Z., 285
Jewett, D.C., Schaal, D.W., Cleary, J., Thompson, T. and Levine, A.S., The discriminative stimulus effects of neuropeptide Y, 165

Kamada, ., see Kitagawa, K., 203
Katayama, Y., see Yoshino, A., 106
Kawamata, T., see Yoshino, A., 106
Kellogg, C.K., see Bitran, D., 157
Kellogg, C.K., see Primus, R.J., 299
Kelly, M.E., see McIntyre, D.C., 279
Kikuchi, H., see Akiyama, Y., 120
Kimura, K., see Kitagawa, K., 203
Kirkman, J.A., see Duncan, G.E., 84
Kitagawa, K., Matsumoto, M., Kuwabara, K., Tagaya, M., Ohtsuki, T., Hata, R., Ueda, H., Handa, N., Kimura, K., and Kamada, T., 'Ischemic tolerance' phenomenon detected in various brain regions, 203
Klee, M.R., see Shirasaki, T., 77
Koh, J.-y., Palmer, E., Lin, A. and Cotman C.W., A metabotropic glutamate receptor agonist does not mediate neuronal degeneration in cortical culture 338
Koplas, P.A., see Duncan, G.E., 84
Koshimura, K., see Akiyama, Y., 120
Krause, T.L., Marquis, R.E., Lyckman, A.W., Ballinger, M.L. and Bittner, G.D., Rapid artificial restoration of electrical continuity across a crush lesion of a giant axon, 350
Kurachi, M., see Minabe, Y., 162
Kuwabara, K., see Kitagawa, K., 203

LaMotte, R.H., see Petersen, M., 20
Lehmenkühler, A., Nicholson, C. and Speckmann, E.-J., Threshold extracellular concentration distribution of penicillin for generation of epileptic focus measured by diffusion analysis, 292
Leslie, M., Forger, N.G. and Breedlove, S.M., Sexual dimorphism and androgen effects on spinal motoneurons innervating the rat flexor digitorum brevis, 269
Levine, A.S., see Jewett, D.C., 165
Levy, W.B., see Tomasulo, R.A., 27
Lewis, M.H., see Mileson, B.E., 1
Lewis, S.J., see Meller, S.T., 61

Brain Research, 561 (1991) 358–359
Elsevier

Author Index

Akaike, N., see Shirasaki, T., 77
Akiyama, Y., Ito, A., Koshimura, K., Ohue, T., Yamagata, S., Miwa, S. and Kikuchi, H., Effects of transient forebrain ischemia and reperfusion on function of dopaminergic neurons and dopamine reuptake in vivo in rat striatum, 120
Appenteng, K., see Saha, S., 128
Arnold, A.P., see Schlinger, B.A., 99

Ballinger, M.L., see Krause, T.L., 350
Batten, T.F.C., see Saha, S., 128
Becker, C.-M., see Naas, E., 139
Becker, D.P., see Yoshino, A., 106
Beitner-Johnson, D., Guitart, X. and Nestler, E.J., Dopaminergic brain-reward regions of Lewis and Fischer rats display different levels of tyrosine hydroxylase and other morphine- and cocaine-regulated phosphoproteins, 147
Betz, H., see Naas, E., 139
Bicknell, R.J., see Oldfield, B.J., 151
Bilkey, D.K., see Yamada, N., 324
Bitran, D., Hilvers, R.J. and Kellogg, C.K., Anxiolytic effects of 3α-hydroxy-5α[β]-pregnan-20-one: endogenous metabolites of progesterone that are active at the $GABA_A$ receptor, 157
Bittner, G.D., see Krause, T.L., 350
Bohn, M.C., see Cunningham, L.A., 192
Breedlove, S.M., see Leslie, M., 269
Breese, G.R., see Duncan, G.E., 84
Bridges, R.J., Hatalski, C., Shim, S.N. and Nunn, P.B., Gliotoxic properties of the *Lathyrus* excitotoxin β-*N*-oxalyl-L-α,β-diaminopropionic acid (β-L-ODAP), 262
Brody, M.J., see Meller, S.T., 61

Campbell, J.N., see Meyer, R.A., 252
Cerne, R., see Gerber, R., 236
Chalmers, D.T. and Watson, S.J., Comparative anatomical distribution of 5-HT_{1A} receptor mRNA and 5-HT_{1A} binding in rat brain — a combined in situ hybridisation/in vitro receptor autoradiographic study, 51
Chapin, J.K., see Nicolelis, M.A.L., 344
Chiu, A.T., see DeNoble, V.J., 230
Cleary, J., see Jewett, D.C., 165
Cler, J.A., see Rao, T.S., 43
Cohen, R.H., see Meyer, R.A., 252
Connor, J.A., see Hyndman, A.G., 318
Contreras, P.C., see Rao, T.S., 43
Cotman, C.W., see Koh, J.-y., 338
Crawford, C., see Iwamoto, N., 177
Cunningham, L.A., Hansen, J.T., Short, M.P. and Bohn, M.C., The use of genetically altered astrocytes to provide nerve growth factor to adrenal chromaffin cells grafted into the striatum, 192

Davis, K.D., see Meyer, R.A., 252
Deng, M.H., Lynch, H.J. and Wurtman, R.J., Melatonin and its precursors in Y79 human retinoblastoma cells: effect of sodium butyrate, 274
DeNoble, K.F., see DeNoble, V.J., 230
DeNoble, V.J., DeNoble, K.F., Spencer, K.R., Chiu, A.T., Wong, P.C. and Timmermans, P.B.M.W.M., Nonpeptide angiotensin II receptor antagonist and angiotensin-converting enzyme inhibitor: effect on a renin-induced deficit of a passive avoidance response in rats, 230
Dickson, K.L., see Seckl, J.R., 332
Dilworth, V.M., see Rao, T.S., 43
Dufresne, C., see McIntyre, D.C., 279
Duncan, G.E., Little, K.Y., Koplas, P.A., Kirkman, J.A., Breese, G.R. and Stumpf, W.E., β-Adrenergic receptor distribution in human and rat hippocampal formation: marked species differences, 84

Emmett, M.R., see Rao, T.S., 43
Emori, K., see Minabe, Y., 162
Emson, P.C., see Iwamoto, N., 177

Farley, L.G., see Silver, W.L., 212
Finger, T.E., see Silver, W.L., 212
Fink, G., see Seckl, J.R., 332
Forger, N.G., see Leslie, M., 269
Frusciante, M.J., see Nussenzveig, I.Z., 285

Gebhart, G.F., see Meller, S.T., 61
Gerber, G., Cerne, R. and Randić, M., Participation of excitatory amino acid receptors in the slow excitatory synaptic transmission in rat spinal dorsal horn, 236
Gibson, M.J., see Saitoh, Y., 11
Gnahn, H., see Naas, E., 139
Gray, N.M., see Rao, T.S., 43
Guitart, X., see Beitner-Johnson, D., 147

Haji, A., see Takeda, R., 307
Handa, N., see Kitagawa, K., 203
Hansen, J.T., see Cunningham, L.A., 192
Hata, R., see Kitagawa, K., 203
Hatalski, C., see Bridges, R.J., 262
Hilvers, R.J., see Bitran, D., 157
Hockberger, P.E., see Hyndman, A.G., 318
Hoebel, B.G., see Rada, P., 354
Horie, H., Pamplin, P.J. and Yokota, T., Inhibition of nociceptive neurons in the shell region of nucleus ventralis posterolateralis following conditioning stimulation of the periaqueductal grey of the cat. Evidence for an ascending inhibitory pathway, 35
Hovda, D.A., see Yoshino, A., 106
Hutchins, B., Evidence for a direct retinal projection to the anterior pretectal nucleus in the cat, 169
Hyndman, A.G., Hockberger, P.E., Zeevalk, G.D. and Connor, J.A., Transferrin can alter physiological properties of retinal neurons, 318

Ito, A., see Akiyama, Y., 120
Iwamoto, N., Thangnipon, W., Crawford, C. and Emson, P.C., Localization of calpain immunoreactivity in senile plaques and in neurones undergoing neurofibrillary degeneration in Alzheimer's disease, 177
Iyengar, S., see Rao, T.S., 43

Javitt, D.C., see Nussenzveig, I.Z., 285
Jewett, D.C., Schaal, D.W., Cleary, J., Thompson, T. and Levine, A.S., The discriminative stimulus effects of neuropeptide Y, 165

Kamada, ., see Kitagawa, K., 203
Katayama, Y., see Yoshino, A., 106
Kawamata, T., see Yoshino, A., 106
Kellogg, C.K., see Bitran, D., 157
Kellogg, C.K., see Primus, R.J., 299
Kelly, M.E., see McIntyre, D.C., 279
Kikuchi, H., see Akiyama, Y., 120
Kimura, K., see Kitagawa, K., 203
Kirkman, J.A., see Duncan, G.E., 84
Kitagawa, K., Matsumoto, M., Kuwabara, K., Tagaya, M., Ohtsuki, T., Hata, R., Ueda, H., Handa, N., Kimura, K., and Kamada, T., 'Ischemic tolerance' phenomenon detected in various brain regions, 203
Klee, M.R., see Shirasaki, T., 77
Koh, J.-y., Palmer, E., Lin, A. and Cotman C.W., A metabotropic glutamate receptor agonist does not mediate neuronal degeneration in cortical culture 338
Koplas, P.A., see Duncan, G.E., 84
Koshimura, K., see Akiyama, Y., 120
Krause, T.L., Marquis, R.E., Lyckman, A.W., Ballinger, M.L. and Bittner, G.D., Rapid artificial restoration of electrical continuity across a crush lesion of a giant axon, 350
Kurachi, M., see Minabe, Y., 162
Kuwabara, K., see Kitagawa, K., 203

LaMotte, R.H., see Petersen, M., 20
Lehmenkühler, A., Nicholson, C. and Speckmann, E.-J., Threshold extracellular concentration distribution of penicillin for generation of epileptic focus measured by diffusion analysis, 292
Leslie, M., Forger, N.G. and Breedlove, S.M., Sexual dimorphism and androgen effects on spinal motoneurons innervating the rat flexor digitorum brevis, 269
Levine, A.S., see Jewett, D.C., 165
Levy, W.B., see Tomasulo, R.A., 27
Lewis, M.H., see Mileson, B.E., 1
Lewis, S.J., see Meller, S.T., 61

Lin, A., see Koh, J.-y., 338
Lin, R.C.S., see Nicolelis, M.A.L., 344
Little, K.Y., see Duncan, G.E., 84
Lyckman, A.W., see Krause, T.L., 350
Lynch, H.J., see Deng, M.H., 274

Mailman, R.B., see Mileson, B.E., 1
Mark, G.P., see Rada, P., 354
Marquis, R.E., see Krause, T.L., 350
Matsumoto, M., see Kitagawa, K., 203
McAllen, R.M., see Oldfield, B.J., 151
McIntyre, D.C., Kelly, M.E. and Dufresne, C., Suppression of amygdala kindling with massed stimulation: effect of noradrenaline antagonists, 279
McKinley, M.J., see Oldfield, B.J., 151
Meller, S.T., Lewis, S.J., Brody, M.J. and Gebhart, G.F., The peripheral nociceptive actions of intravenously administered 5-HT in the rat requires dual activation of both 5-HT_2 and 5-HT_3 receptor subtypes, 61
Meyer, R.A., Davis, K.D., Cohen, R.H., Treede, R.-D. and Campbell, J.N., Mechanically insensitive afferents (MIAs) in cutaneous nerves of monkey, 252
Mick, S.J., see Rao, T.S., 43
Mileson, B.E., Lewis, M.H. and Mailman, R.B., Dopamine receptor 'supersensitivity' occurring without receptor up-regulation, 1
Minabe, Y., Emori, K. and Kurachi, M., Two types of neuroplasticities in the kindling phenomenon, 162
Miwa, S., see Akiyama, Y., 120

Naas, E., Zilles, K., Gnahn, H., Betz, H., Becker, C.-M. and Schröder, H., Glycine receptor immunoreactivity in rat and human cerebral cortex, 139
Nakaye, T., see Shirasaki, T., 77
Nestler, E.J., see Beitner-Johnson, D., 147
Nicholson, C., see Lehmenkühler, A., 292
Nicolelis, M.A.L., Chapin, J.K. and Lin, R.C.S., Thalamic plasticity induced by early whisker removal in rats, 344
Nunn, P.B., see Bridges, R.J., 262
Nussenzveig, I.Z., Sircar, R., Wong, M.-L., Frusciante, M.J., Javitt, D.C. and Zukin, S.R., Polyamine effects upon *N*-methyl-D-aspartate receptor functioning: differential alteration by glutamate and glycine site antagonists, 285

Ohtsuki, T., see Kitagawa, K., 203
Ohue, T., see Akiyama, Y., 120
Oldfield, B.J., Bicknell, R.J., McAllen, R.M., Weisinger, R.S. and McKinley, M.J., Intravenous hypertonic saline induces Fos immunoreactivity in neurons throughout the lamina terminalis, 151

Palmer, E., see Koh, J.-y., 338
Pamplin, P.J., see Horie, H., 35
Paton, J.F.R., Rogers, W.T. and Schwaber, J.S., The ventrolateral medulla as a source of synaptic drive to rhythmically firing neurons in the cardiovascular nucleus tractus solitarius of the rat, 217
Pessah, I.N., see Zimanyi, I., 181
Petersen, M. and LaMotte, R.H., Relationships between capsaicin sensitivity of mammlian sensory neurons, cell size and type of voltage gated Ca-currents, 20
Pothos, E., see Rada, P., 354
Primus, R.J. and Kellogg, C.K., Gonadal status and pubertal age influence the responsiveness of the benzodiazepine/GABA receptor complex to environmental challenge in male rats, 299

Rada, P., Pothos, E., Mark, G.P. and Hoebel, B.G., Microdialysis evidence that acetylcholine in the nucleus accumbens is involved in morphine withdrawal and its treatment with clonidine, 354
Randić, M., see Gerber, G., 236
Rao, T.S., Mick, S.J., Cler, J.A., Emmett, M.R., Dilworth, V.M., Contreras, P.C., Gray, N.M., Wood, P.L.and Iyengar, S., Effects of sigma ligands on mouse cerebellar cyclic guanosine monophosphate (cGMP) levels in vivo: further evidence for a functional modulation of *N*-methyl-D-aspartate (NMDA) receptor complex-mediated events by sigma ligands, 43
Rogers, W.T., see Paton, J.F.R., 217

Saha, S., Appenteng, K. and Batten, T.F.C., Quantitative analysis and postsynaptic targets of GABA-immunoreactive boutons within the rat trigeminal motor nucleus, 128
Saitoh, Y., Silverman, A.-J. and Gibson, M.J., Norepinephrine neurons in mouse locus coeruleus express c-*fos* protein after *N*-methyl-D,L-aspartic acid (NMDA) treatment: relation to LH release, 11
Schaal, D.W., see Jewett, D.C., 164
Schlinger, B.A. and Arnold, A.P., Androgen effects on the development of the zebra finch song system, 99
Schröder, H., see Naas, E., 139
Schwaber, J.S., see Paton, J.F.R., 217
Seckl, J.R., Dickson, K.L., Yates, C. and Fink, G., Distribution of glucocorticoid and mineralocorticoid receptor messenger RNA expression in human postmortem hippocampus, 332
Shim, S.N., see Bridges, R.J., 262
Shirasaki, T., Klee, M.R., Nakaye, T. and Akaike, N., Differential blockade of bicuculline and strychnine on GABA- and glycine-induced responses in dissociated rat hippocampal pyramidal cells, 77
Short, M.P., see Cunningham, L.A., 192
Silver, W.L., Farley, L.G. and Finger, T.E., The effects of neonatal capsaicin administration on trigeminal nerve chemoreceptors in the rat nasal cavity, 212
Silverman, A.-J., see Saitoh, Y., 11
Sircar, R., see Nussenzveig, I.Z., 285
Speckmann, E.-J., see Lehmenkühler, A., 292
Spencer4, K.R., see DeNoble, V.J., 230
Steward, O., see Tomasulo, R.A., 27
Stumpf, W.E., see Duncan, G.E., 84

Tagaya, M., see Kitagawa, K., 203
Takeda, R. and Haji, A., Synaptic response of bulbar respiratory neurons to hypercapnic stimulation in peripherally chemodenervated cats, 307
Thangnipon, W., see Iwamoto, N., 177
Thompson, T., see Jewett, D.C., 165
Thornhill, R.A., Cholinergic nicotenic receptors in the vestibular epithelia, 174
Tietz, E.I., see Xie, X.-H., 69
Timmermans, P.B.M.W.M., see DeNoble, V.J., 230
Tomasulo, R.A., Levy, W.B. and Steward, O., LTP-associated EPSP/spike dissociation in the dentate gyrus: GABAergic and non-GABAergic components, 27
Treede, R.-D., see Meyer, R.A., 252

Ueda, H., see Kitagawa, K., 203

Von Euler, G., Biochemical characterization of the intramembrane interaction between neurotensin and dopamine D_2 receptors in the rat brain, 93

Watson, S.J., see Chalmers, D.T., 51
Weisinger, R.S., see Oldfield, B.J., 151
Wong, M.-L., see Nussenzveig, I.Z., 285
Wong, P.C., see DeNoble, V.J., 230
Wood, P.L., see Rao, T.S., 43
Wurtman, R.J., see Deng, M.H., 274

Xie, X.-H. and Tietz, E.I., Chronic benzodiazepine treatment of rats induces reduction of paired-pulse inhibition in CA1 region of in vitro hippocampus, 69

Yamada, N. and Bilkey, D.K., Kindling-induced persistent alterations in the membrane and synaptic properties of CA1 pyramidal neurons, 324
Yamagata, S., see Akiyama, Y., 120
Yates, C., see Seckl, J.R., 332
Yokota, T., see Horie, H., 35
Yoshino, A., Hovda, D.A., Kawamata, T., Katayama, Y. and Becker, D.P., Dynamic changes in local cerebral glucose utilization following cerebral concussion in rats: evidence of a hyper- and subsequent hypometabolic state, 106

Zeevalk, G.D., see Hyndman, A.G., 318
Zilles, K., see Naas, E., 139
Zimanyi, I. and Pessah, I.N., Pharmacological characterization of the specific binding of [^{3}H]ryanodine to rat brain microsomal membranes 181
Zukin, S.R., see Nussenzveig, I.Z., 285